

October 2018

OCCURRENCE OF ESCHERICHIA COLI O157 IN GROUND MEATCOMMERCIALIZED IN LEBANESE URBAN CITIES

MICHEL AFRAM Lebanese Agricultural Research Institute, Lebanon
Lebanese Agricultural Research Institute, Lebanon, lari@lari.gov.lb

JOSEPH TOUMA Lebanese Agricultural Research Institute, Lebanon
Lebanese Agricultural Research Institute, Lebanon, toumajoseph@hotmail.com

NADA EI-DARRA Assistant Professor, Department of Nutrition & Dietetics, Faculty of Health Sciences,
Beirut Arab University, Lebanon, n.aldarra@bau.edu.lb

Follow this and additional works at: <https://digitalcommons.bau.edu.lb/hwbjournal>



Part of the [Architecture Commons](#), [Business Commons](#), [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

AFRAM, MICHEL Lebanese Agricultural Research Institute, Lebanon; TOUMA, JOSEPH Lebanese Agricultural Research Institute, Lebanon; and EI-DARRA, NADA Assistant Professor, Department of Nutrition & Dietetics, Faculty of Health Sciences, (2018) "OCCURRENCE OF ESCHERICHIA COLI O157 IN GROUND MEATCOMMERCIALIZED IN LEBANESE URBAN CITIES," *BAU Journal - Health and Wellbeing*: Vol. 1 : Iss. 3 , Article 38.

Available at: <https://digitalcommons.bau.edu.lb/hwbjournal/vol1/iss3/38>

This Article is brought to you for free and open access by Digital Commons @ BAU. It has been accepted for inclusion in BAU Journal - Health and Wellbeing by an authorized editor of Digital Commons @ BAU. For more information, please contact ibtihal@bau.edu.lb.

OCCURRENCE OF ESCHERICHIA COLI O157 IN GROUND MEAT COMMERCIALIZED IN LEBANESE URBAN CITIES

Abstract

Escherichia coli O157:H7 (E. coli O157:H7) has been recognized as a major cause of diarrhea and hemolytic-uremic syndrome (HS). The consumption of raw or undercooked meat of bovine origin has been the most common mean of transmitting this organism. No screening of E. coli has been performed in the ground meat commercialized in the Lebanese market. Therefore, the purpose of this study is to evaluate the prevalence of E. coli serotype O157:H7 recovered from ground fresh raw meat collected from different butcheries in Lebanese Urban cities. A total of 73 samples of fresh ground meat was collected from Sin el fil area (n=23), Hadath (n=16), Ghoubayri (n=16) and Antelias (n=18), and investigated for the presence of E. coli O157:H7. E. coli O157:H7 was isolated in 16 (22%) out of 73 meat samples examined. This high prevalence was variable between the different cities, with the highest one in Ghoubayri (43%), followed by Sin el fil (21%), Hadath (12%) and Antelias (11%). This study revealed the presence of E. coli O157:H7 in retail raw meats reaching consumers, especially in crowded urban city such as Ghoubayri. This result is an indication of the poor hygienic level in the different butcheries localized in the Lebanese urban cities, thus reflecting possible risks of infection to people through the consumption of fresh raw/under-cooked meat.

Keywords

Occurrence, E. coli, meat, Lebanon, Food safety

OCCURRENCE OF ESCHERICHIA COLI O157 IN GROUND MEAT COMMERCIALIZED IN LEBANESE URBAN CITIES

MICHEL AFRAM¹, JOSEPH TOUMA², and
NADA EI-DARRA³

¹ Lebanese Agricultural Research Institute, Lebanon

² Lebanese Agricultural Research Institute, Lebanon

³ Assistant Professor, Department of Nutrition & Dietetics, Faculty of Health Sciences,
Beirut Arab University, Lebanon

ABSTRACT: *Escherichia coli O157:H7 (E. coli O157:H7) has been recognized as a major cause of diarrhea and hemolytic- uremic syndrome (HS). The consumption of raw or undercooked meat of bovine origin has been the most common mean of transmitting this organism. No screening of E.coli has been performed in the ground meat commercialized in the Lebanese market. Therefore, the purpose of this study is to evaluate the prevalence of E. coli serotype O157:H7 recovered from ground fresh raw meat collected from different butchereries in Lebanese Urban cities. A total of 73 samples of fresh ground meat was collected from Sin el fil area (n=23), Hadath (n=16), Ghoubayri (n=16) and Antelias (n=18), and investigated for the presence of E. coli O157: H7. E. coli O157:H7 was isolated in 16 (22%) out of 73 meat samples examined. This high prevalence was variable between the different cities, with the highest one in Ghoubayri (43%), followed by Sin el fil (21%), Hadath (12%) and Antelias (11%). This study revealed the presence of E. coli O157:H7 in retail raw meats reaching consumers, especially in crowded urban city such as Ghoubayri. This result is an indication of the poor hygienic level in the different butchereries localized in the Lebanese urban cities, thus reflecting possible risks of infection to people through the consumption of fresh raw/under-cooked meat.*

1. INTRODUCTION

Nowadays, Food borne outbreaks represent a major health problem especially in developing countries such as Lebanon, where food safety scandals are a serious concern. Foodborne outbreaks are of increasing occurrence in Lebanon and in response to that, the Ministry of Public Health have done a series of inspections and publicized a list of restaurants, supermarkets and other food provider services that did not meet the ministry's food regulation standards (Institute of Health Management and Social Protection, USJ, WHO, & MoPH, 2012).

Different factors might increase the food-borne disease outbreak origin challenging, e.g. population growth, changing eating habits, globalization of food supply chains, production and processing innovations, and microbiological adaptation (Newell et al., 2010).

Literature studies have shown that 75% of food borne illness outbreaks is related to improper food handling practices in food establishments (Gizaw, Gebrehiwot, & Teka, 2014). Poor hygiene status among food handlers is being considered as one of the major risk factors in the transmission of bacteria and contamination of food thus posing serious health risks to consumers (Scallan et al., 2011). Consequently, food safety training should be implemented on food handlers to increase awareness about the proper hygienic practices (Tan, Lee, & Mahyudin, 2014).

Shiga-toxin producing *Escherichia coli* are currently considered as important emerging food-borne bacterial pathogens of public health concern. They are causing sporadic and epidemic human cases of hemorrhagic colitis (HC), which can lead to severe and life threatening hemolytic and uremic syndrome (HUS), especially in young infants (Madic, 2010). The majority of the cases are due to the ingestion of food or water contaminated with STEC strains belonging to the serotype O157:H7. The Centers for Disease Control and Prevention reported *Escherichia coli* as one of the three most common food-borne infections (CDC., 2008). Traditional methods for the detection of *E. coli* O157 use enrichment, followed by isolation on Sorbitol MacConkey Agar. The methods are time-consuming, and competing flora may obscure O157 colonies thereby giving rise to false negative results. Recently, many companies have developed rapid

methods for detection that are specific, faster and more sensitive than traditional culture methods. Enzyme-linked immunosorbent fluorescent assays (VIDAS-UP) is one of these methods (HILDA NYATI, ANNET HEUVELINK & ZWARTKRUIS, 2012).

Since *Escherichia coli* (*E. coli*) is being considered as one of the major causes in food borne outbreaks in Lebanon (Harakeh et al., 2005), we aimed in this study at screening the prevalence of *E. coli* serotype O157:H7 recovered from ground fresh raw meat collected from different butcheries in Lebanese Urban cities Sin el fil area, Hadath, Ghoubayri, and Antelias. The samples will be tested by a rapid method, followed by a confirmation with the reference method of *E. coli* identification (ISO EN-16654)

2. MATERIALS AND METHODS

2.1 Food outbreak prevalence

A screening was conducted with the Ministry of Public Health to check the prevalence of foodborne outbreaks among Lebanese hospitals between January 2014 till January 2015. length of paper title may not exceed 80 characters. Paper title is typed bold uppercase letters, 14 pt., and placed centered. Author(s) name(s) should be typed in plain capital letters, size 12, not bold, centered, and placed one space under the paper title with footnote giving the author(s) title(s) and affiliation(s).

2.2 Sample size and sampling procedure

A total of 73 samples of fresh ground meat were collected to test the prevalence of *E. coli* serotype O157:H7 recovered from ground fresh raw meat collected from different butcheries in Lebanese Urban cities. The samples were randomly collected from four Lebanese urban cities, Sin el fil area (n=23), Hadath (n=16), Ghoubayri (n=16) and Antelias (n=18). The samples were stored at a temperature 1-4°C, the subjected to the analysis within 24 hours.

2.3 Microbiological procedure

The fresh ground meat samples were subjected to a rapid screening, VIDAS UP, for the detection of *E. coli* O157. Afterwards, the samples were subjected to the reference method of *E. coli* O157 identification, which is the ISO-EN-16654.

3. RESULTS AND DISCUSSION

3.1 Food-borne Outbreaks in Lebanon

Captions In collaboration with the ministry of Public Health, a screening of the Foodborne outbreak cases among the hospitals in Lebanon was conducted between January 2014 and 2015 (Table 1). During this period, 109 foodborne outbreak cases were found. In 2014, 39 cases were identified, with 10 out of 39 were mainly due to contaminated chicken, 8 were related to the raw meat. This number almost doubled in 2015 to reach 70 cases in total, with 20 due to chicken and 15 due to raw meat. Chicken and raw meat are potential vehicles for transmitting food-borne diseases (Zhao et al., 2001). This screening is important since focusing on foods that are prominent in outbreaks will better guide the food industries and governments in targeting interventions (Crowe, Mahon, Vieira, & Gould, 2015). Consequently, there is a need to increase implementation of hazard analysis of critical control point (HACCP) in meat industries and consumer food safety education efforts.

Table 1. Prevalence of Foodborne outbreaks in Lebanon between January 2014 and 2015

Vehicle category	Number of outbreaks, 2014	Number of outbreaks, 2015
Chicken	10	20
Meat	3	7
Raw meat	8	15
Mayonnaise	2	5
Egg dishes	2	3
Cheese	1	1

Fish	1	3
Dessert	4	4
Arab sweets	2	5
Milk	1	0
Mixed vehicle	5	7
Total	39	70

3.2 Identification of pathogenic organisms in the food causing outbreaks

C After checking the main food source of the outbreaks, the identification of the involved pathogenic organisms becomes highly crucial for surveillance, prevention, and control of food-borne diseases.

Table 2 showed that *Escherichia coli* O157:H7, *Staphylococcus aureus*, *Listeria monocytogenes* and *Salmonella* spp. were detected in all the food samples (466), which is in correlation with the literature findings that show that these microorganisms are able to survive as potential food pathogens (Park, Lee, & Kim, 2006). The results showed that *Escherichia coli* is almost responsible of half of the detected contamination of food samples. Since most outbreaks with *E.coli* O157:H7 infection have been linked to foods of bovine origin (Abu-Ali et al., 2009), we decide to investigate further fresh ground meat from different locations located in Lebanese urban cities for their contamination with *E.coli* O157:H7.

Table 2. Microorganisms isolated from food causing foodborne outbreaks

Isolated microorganisms	Number	%
<i>Escherichia coli</i>	205	44
<i>Staphylococcus aureus</i>	107	23
<i>Listeria monocytogenes</i>	19	4
<i>Salmonella</i>	135	29
<i>Total</i>	466	100

3.3 *E.coli* occurrence in fresh ground meat located in different Lebanese Urban Cities

C Table 3 represents the *E.coli* serotype O157:H7 occurrence in fresh ground meat (n=73) located in different Lebanese Urban Cities, Sin el fil area (n=23), Hadath (n=16), Ghoubayri (n=16) and Antelias (n=18). *E. coli* O157:H7 was isolated in 16 (22%) out of 73 meat samples examined, which is in accordance with the literature data that showed that ground beef has been the food most of ten associated with *E. coli* O157 (Ateba & Bezuidenhout, 2008). This contamination could be due to the slaughtering process of the cattle which may contaminate the meat with the *E. coli* bacteria from the cattle intestine. Studies on cattle have shown that 2.8- 1000/3.3cattle carry *E. Coli* O157:H7. This latter has shown too survive in the cattle feces for up to 70 days (Wang, Zhao, & Doyle, 1996). The high prevalence was variable between the different cities, with the highest one in Ghoubayri (43%), followed by Sin el fil (21%), Hadath (12%) and Antelias (11%). This study confirmed the presence of *E. coli* O157:H7 in retail raw meats reaching consumers, especially in crowded urban city such as Ghoubayri. This result is an indication of the poor hygienic level in the different butcheries localized in the Lebanese urban cities, thus reflecting possible risks of infection to people through the consumption of fresh raw/under-cooked meat.

Table 3. E.coli occurrence in fresh ground meat located in different Lebanese Urban Cities

Samples (73)	%
Sin el fil area (n=23),	21
Hadath (n=16),	12
Ghoubayri (n=16)	43
Antelias (n=18)	11

4. CONCLUSIONS

The Surveillance for Foodborne Disease Outbreaks in Lebanon, suggests that chicken and meat has caused the most food-related sickness among the public. E. coli is the main pathogen contaminating meat during slaughter and processing. This study pointed to the high prevalence of E. coli in the meat distributed in many Lebanese Urban cities and suggest the need for proper control strategies for the prevention of contaminated animal products. Effective measures to promote safe food-handling practices could reduce poultry-associated outbreaks and illnesses.

ACKNOWLEDGEMENT

The corresponding author thankfully acknowledges the Lebanese Agricultural Research Institute for its continuous support.

REFERENCES

- Abu-Ali, G. S., Lacher, D. W., Wick, L. M., Qi, W., Whittam, T. S., Donnenberg, M., ... Felsenstein, J. (2009). Genomic diversity of pathogenic *Escherichia coli* of the EHEC 2 clonal complex. *BMC Genomics*, 10(1), 296. <http://doi.org/10.1186/1471-2164-10-296>
- Ateba, C. N., & Bezuidenhout, C. C. (2008). Characterisation of *Escherichia coli* O157 strains from humans, cattle and pigs in the North-West Province, South Africa. *International Journal of Food Microbiology*. <http://doi.org/10.1016/j.ijfoodmicro.2008.08.011>
- CDC. (2008). *Escherichia coli*. Centres for Disease Control and Prevention.
- Crowe, S. J., Mahon, B. E., Vieira, A. R., & Gould, L. H. (2015). Vital Signs: Multistate Foodborne Outbreaks — United States, 2010–2014. *MMWR. Morbidity and Mortality Weekly Report*, 64(43), 1221–1225. <http://doi.org/10.15585/mmwr.mm6443a4>
- Gizaw, Gebrehiwot, & Teka. (2014). Food Safety Practice and Associated Factors of Food Handlers Working in Substandard Food Establishments in Gondar Town, Northwest Ethiopia, 2013/14. *International Journal of Food Science, Nutrition and Dietetics*, 138–146. <http://doi.org/10.19070/2326-3350-1400027>
- Harakeh, S., Yassine, H., Gharios, M., Barbour, E., Hajjar, S., El-Fadel, M., ... Tannous, R. (2005). Isolation, molecular characterization and antimicrobial resistance patterns of *Salmonella* and *Escherichia coli* isolates from meat-based fast food in Lebanon. *Science of the Total Environment*, 341(1–3), 33–44. <http://doi.org/10.1016/j.scitotenv.2004.09.025>
- HILDA NYATI, ANNET HEUVELINK, C. V. H., & ZWARTKRUIS. (2012). Evaluation of test-kits for the detection of *Escherichia coli* O157 in raw meats and cattle faeces. *International Journal of Food Studies*, 1, 126–134.
- Institute of Health Management and Social Protection, USJ, WHO, & MoPH. (2012). National Health Statistics Report in Lebanon, 165.
- Madic, J. (2010). Methods for detection of Shiga-Toxin producing *Escherichia coli* (STEC). *NATO Science for Peace and Security Series A: Chemistry and Biology*. <http://doi.org/10.1007/978-90-481-8544-3-4>
- Newell, D. G., Koopmans, M., Verhoef, L., Duizer, E., Aidara-Kane, A., Sprong, H., ... Kruse, H. (2010). Food-borne diseases - The challenges of 20years ago still persist while new ones continue to emerge. *International Journal of Food Microbiology*, 139(SUPPL. 1). <http://doi.org/10.1016/j.ijfoodmicro.2010.01.021>
- Park, Y. S., Lee, S. R., & Kim, Y. G. (2006). Detection of *Escherichia coli* O157:H7, *Salmonella* spp., *Staphylococcus aureus* and *Listeria monocytogenes* in kimchi by multiplex polymerase chain reaction

- (mPCR). *Journal of Microbiology (Seoul, Korea)*, 44(1), 92–7. <http://doi.org/2331>
- Scallan, E., Hoekstra, R. M., Angulo, F. J., Tauxe, R. V., Widdowson, M. A., Roy, S. L., ... Griffin, P. M. (2011). Foodborne illness acquired in the United States-Major pathogens. *Emerging Infectious Diseases*, 17(1), 7–15. <http://doi.org/10.3201/eid1701.P11101>
 - Tan, S. L., Lee, H. Y., & Mahyudin, N. a. (2014). Antimicrobial resistance of Escherichia coli and Staphylococcus aureus isolated from food handler's hands. *Food Control*, 44, 203–207. <http://doi.org/10.1016/j.foodcont.2014.04.008>
 - Wang, G., Zhao, T., & Doyle, M. P. (1996). Fate of enterohemorrhagic Escherichia coli O157:H7 in bovine feces. *Applied and Environmental Microbiology*, 62(7), 2567–2570.
 - Zhao, C., Ge, B., De Villena, J., Sudler, R., Yeh, E., Zhao, S., ... Meng, J. (2001). Prevalence of Campylobacter spp., Escherichia coli, and Salmonella Serovars in Retail Chicken, Turkey, Pork, and Beef from the Greater Washington, D.C., Area. *Applied and Environmental Microbiology*, 67(12), 5431–5436. <http://doi.org/10.1128/AEM.67.12.5431-5436.2001>